

PCT

WORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau




INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification 6 : C07D 401/06, 413/06, 211/60, A61K 31/55, 31/445		A1	(11) International Publication Number: WO 96/31497 (43) International Publication Date: 10 October 1996 (10.10.96)
(21) International Application Number: PCT/DK96/00138 (22) International Filing Date: 1 April 1996 (01.04.96) (30) Priority Data: 0403/95 7 April 1995 (07.04.95) DK 1006/95 11 September 1995 (11.09.95) DK (71) Applicant: NOVO NORDISK A/S [DK/DK]; Novo Allé, DK-2880 Bagsværd (DK). (72) Inventors: JØRGENSEN, Tine, Krogh; Stavnsbjerg Allé 80, DK-2730 Herlev (DK). ANDERSEN, Knud, Erik; Nøddehøvdene 122, DK-2765 Smørum (DK). ANDERSEN, Henrik, Sune; Kastelsvej 24 st. th, DK-2100 København Ø (DK). HOHLWEG, Rolf; Nybovej 6, DK-3490 Kvistgaard (DK). MADSEN, Peter; Ulvebjerg 7, DK-2880 Bagsværd (DK). OLSEN, Uffe, Bang; Horsbred 111, DK-2625 Vallensbæk (DK).			(81) Designated States: AL, AM, AT, AU, AZ, BB, BG, BR, BY, CA, CH, CN, CZ, DE, DK, EE, ES, FI, GB, GE, HU, IS, JP, KE, KG, KP, KR, KZ, LK, LR, LS, LT, LU, LV, MD, MG, MK, MN, MW, MX, NO, NZ, PL, PT, RO, RU, SD, SE, SG, SI, SK, TJ, TM, TR, TT, UA, UG, UZ, VN, ARIPO patent (KE, LS, MW, SD, SZ, UG), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG). Published With international search report. Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.

(54) Title: NOVEL HETEROCYCLIC COMPOUNDS

(57) Abstract

A compound of formula (I). The present invention relates to novel N-substituted azaheterocyclic carboxylic acids and esters thereof in which a substituted alkyl chain forms part of the N-substituent or salts thereof, to methods for their preparation, to compositions containing them, and to their use for the clinical treatment of painful, hyperalgesic and/or inflammatory conditions in which C-fibers play a pathophysiological role by eliciting neurogenic pain or inflammation.



FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AM	Armenia	GB	United Kingdom	MW	Malawi
AT	Austria	GE	Georgia	MX	Mexico
AU	Australia	GN	Guinea	NE	Niger
BB	Barbados	GR	Greece	NL	Netherlands
BE	Belgium	HU	Hungary	NO	Norway
BF	Burkina Faso	IE	Ireland	NZ	New Zealand
BG	Bulgaria	IT	Italy	PL	Poland
BJ	Benin	JP	Japan	PT	Portugal
BR	Brazil	KE	Kenya	RO	Romania
BY	Belarus	KG	Kyrgyzstan	RU	Russian Federation
CA	Canada	KP	Democratic People's Republic of Korea	SD	Sudan
CF	Central African Republic	KR	Republic of Korea	SE	Sweden
CG	Congo	KZ	Kazakhstan	SG	Singapore
CH	Switzerland	LJ	Liechtenstein	SI	Slovenia
CI	Côte d'Ivoire	LK	Sri Lanka	SK	Slovakia
CM	Cameroon	LR	Liberia	SN	Senegal
CN	China	LT	Lithuania	SZ	Swaziland
CS	Czechoslovakia	LU	Luxembourg	TD	Chad
CZ	Czech Republic	LV	Latvia	TG	Togo
DE	Germany	MC	Monaco	TJ	Tajikistan
DK	Denmark	MD	Republic of Moldova	TT	Trinidad and Tobago
EE	Estonia	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	UG	Uganda
FI	Finland	MN	Mongolia	US	United States of America
FR	France	MR	Mauritania	UZ	Uzbekistan
GA	Gabon			VN	Viet Nam

Novel heterocyclic Compounds

5

Field of the Invention

The present invention relates to novel N-substituted azaheterocyclic carboxylic acids and esters thereof in which a substituted alkyl chain forms part of the N-substituent or salts thereof, to methods for their preparation, to compositions containing them, and to their use for the clinical treatment of painful, hyperalgesic and/or inflammatory conditions in which C-fibers play a pathophysiological role by eliciting neurogenic pain or inflammation. The invention also relates to the use of the present compounds for the treatment of insulin resistance in non-insulin-dependent diabetes mellitus (NIDDM) or aging, the present compounds knowing to interfere with neuropeptide containing C-fibres and hence inhibit the secretion and circulation of insulin antagonizing peptides like CGRP or amylin.

10
15Background of the Invention

20

The nervous system exerts a profound effect on the inflammatory response. Antidromic stimulation of sensory nerves results in localized vasodilation and increased vascular permeability (Janecso et al. Br. J. Pharmacol. 1967, 31, 138-151) and a similar response is observed following injection of peptides known to be present in sensory nerves. From this and other data it is postulated that peptides released from sensory nerve endings mediate many inflammatory responses in tissues like skin, joint, urinary tract, eye, meninges, gastro-intestinal and respiratory tracts. Hence inhibition of sensory nerve peptide release and/or activity, may be useful in treatment of, for example arthritis, dermatitis, rhinitis, asthma, cystitis, gingivitis, thrombo-phlebitis, glaucoma, gastro-intestinal diseases or migraine.

25
30

Further, the potent effects of CGRP on skeletal muscle glycogen synthase activity and muscle glucose metabolism, together with the notion that this peptide is released from the neuromuscular junction by nerve excitation, suggest that CGRP may play a physiological role in skeletal muscle glucose metabolism by directing

35

the phosphorylated glucose away from glycogen storage and into the glycolytic and oxidative pathways (Rossetti et al. Am. J. Physiol. 264, E1-E10, 1993). This peptide may represent an important physiological modulator of intracellular glucose trafficking in physiological conditions, such as exercise, and may also
5 contribute to the decreased insulin action and skeletal muscle glycogen synthase in pathophysiological conditions like NIDDM or aging-associated obesity (Melnik et al. Obesity Res. 3, 337-344, 1995) where circulating plasma levels of CGRP are markedly increased. Hence inhibition of release and/or activity of the neuropeptide CGRP may be useful in the treatment of insulin resistance related
10 to type 2 diabetes or aging.


In US Patent No. 4,383,999 and No. 4,514,414 and in EP 236342 as well as in EP 231996 some derivatives of N-(4,4-disubstituted-3-butenyl)azaheterocyclic carboxylic acids are claimed as inhibitors of GABA uptake. In EP 342635 and EP
15 374801, N-substituted azaheterocyclic carboxylic acids in which an oxime ether group and vinyl ether group forms part of the N-substituent respectively are claimed as inhibitors of GABA uptake. Further, in WO 9107389 and WO 9220658, N-substituted azacyclic carboxylic acids are claimed as GABA uptake inhibitors. EP 221572 claims that 1-aryloxyalkylpyridine-3-carboxylic acids are
20 inhibitors of GABA uptake.

Description of the Invention

The present invention relates to novel N-substituted azaheterocyclic carboxylic

3

acids and esters thereof of formula I



wherein R^1 and R^2 independently are hydrogen, halogen, trifluoromethyl,
 5 hydroxy, C_{1-6} -alkyl or C_{1-6} -alkoxy; and

Y is $-\underline{C}H_2-\underline{N}-CH_2-$, $-CH_2-\underline{N}-\underline{C}H_2-$, $-(\underline{C}=\underline{O})-\underline{N}-CH_2-$, $-CH_2-\underline{N}-(\underline{C}=\underline{O})-$,
 $-CH_2-\underline{C}=\underline{C}H-$, $-\underline{C}H=\underline{C}-CH_2-$, $-\underline{C}H_2-\underline{C}H-CH_2-$, $-CH_2-\underline{C}H-\underline{C}H_2-$, $-\underline{C}H_2-\underline{C}=\underline{C}H-$,
 $-\underline{C}H=\underline{C}-\underline{C}H_2-$, $-\underline{O}-\underline{C}H-CH_2-$, $-CH_2-\underline{C}H-\underline{O}-$, $-\underline{S}-\underline{C}H-CH_2-$ or $-CH_2-\underline{C}H-\underline{S}-$ wherein
 only the underscored atoms participate in the ring system; and

10 X is $-O-$, $-S-$, $-C(R^6R^7)-$, $-CH_2CH_2-$, $-CH=CH-CH_2-$, $-CH_2-CH=CH-$, $-CH_2-(C=O)-$,
 $-(C=O)-CH_2-$, $-CH_2CH_2CH_2-$, $-CH=CH-$, $-N(R^8)-(C=O)-$, $-(C=O)-N(R^8)-$, $-O-CH_2-$,
 $-CH_2-O-$, $-S-CH_2-$, $-CH_2-S-$, $-(C=O)-$, $-N(R^9)-$ or $-(S=O)-$ wherein R^6 , R^7 , R^8 and
 R^9 independently are hydrogen or C_{1-6} -alkyl; and
 q is 0 or 1; and

p is 0 or 1; and

r is 1, 2 or 3; and

m is 1 or 2; and

n is 1 when m is 1 and n is 0 when m is 2; and

- 5 R³ and R⁴ each represents hydrogen or may - when m is 2 - together represent a bond; and

R⁵ is -OH or C₁₋₆-alkoxy;

or a pharmaceutically acceptable salt thereof.

- 10 The compounds of formula I may exist as geometric and optical isomers and all isomers and mixtures thereof are included herein. Isomers may be separated by means of standard methods such as chromatographic techniques or fractional crystallization of suitable salts.

- 15 Preferably, the compounds of formula I exist as the individual geometric or optical isomers.

- The compounds according to the invention may optionally exist as pharmaceutically acceptable acid addition salts or - when the carboxylic acid group is
- 20 not esterified - as pharmaceutically acceptable metal salts or - optionally alkylated - ammonium salts.

- Examples of such salts include inorganic and organic acid addition salts such as hydrochloride, hydrobromide, sulphate, phosphate, acetate, fumarate,
- 25 maleate, citrate, lactate, tartrate, oxalate or similar pharmaceutically acceptable inorganic or organic acid addition salts, and include the pharmaceutically acceptable salts listed in Journal of Pharmaceutical Science, 66, 2 (1977)

which are hereby incorporated by reference.

The term "C₁₋₆-alkyl" as used herein, alone or in combination, refers to a straight or branched, saturated hydrocarbon chain having 1 to 6 carbon atoms such as e.g. methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, n-pentyl, 2-methylbutyl, 3-methylbutyl, n-hexyl, 4-methylpentyl, neopentyl, n-hexyl, 1,2-dimethylpropyl, 2,2-dimethylpropyl and 1,2,2-trimethylpropyl.

The term "C₁₋₆-alkoxy" as used herein, alone or in combination, refers to a straight or branched monovalent substituent comprising a C₁₋₆-alkyl group linked through an ether oxygen having its free valence bond from the ether oxygen and having 1 to 6 carbon atoms e.g. methoxy, ethoxy, propoxy, isopropoxy, butoxy, pentoxy.

The term "halogen" means fluorine, chlorine, bromine or iodine.

Illustrative examples of compounds encompassed by the present invention include:

- (R)-1-(3-(6,11-Dioxo-6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;
(R)-1-(3-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;
- (R)-1-(3-(5,11-Dihydro-10H-dibenzo[b,e][1,4]diazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

- (R)-1-(3-(11H-Dibenzo[b,f][1,4]thiazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 5 (R)-1-(3-(11H-Dibenz[b,f][1,4]oxazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;
- (R)-1-(3-(11H-Dibenz[b,f][1,4]oxathiepin-11-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 10 (R)-1-(3-(11H-Dibenzo[b,e][1,4]dithiepin-11-yl)-1-propyl)-3-piperidinecarboxylic acid;
- (R)-1-(3-(11H-Dibenz[b,e][1,4]oxathiepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 15 (R)-1-(3-(11,12-Dihydro-10H-dibenz[b,g][1,5]oxazocin-11-yl)-1-propyl)-3-piperidinecarboxylic acid;
- (R)-1-(3-(11,12-Dihydro-10H-dibenzo[b,g][1,5]thiazocin-11-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 20 1-(3-(11,12-Dihydro-6H-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 25 1-(3-(11,12-Dihydro-5H-dibenzo[a,e]cycloocten-5-ylidene)-1-propyl)-3-piperidinecarboxylic acid;

- 1-(3-(6-Oxo-11,12-dihydro-5H-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 5 1-(3-(7,12-Dihydro-6H-dibenzo[a,d]cycloocten-6-ylidene)-1-propyl)-3-piperidinecarboxylic acid;
- 1-(3-(5-Methyl-5,11-dihydro-dibenz[b,f]azepin-10-ylidene)-1-propyl)-3-piperidinecarboxylic acid;
- 10 1-(3-(6-Oxo-5,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;
- (R)-1-(3-(11-Oxo-10,11-dihydro-5H-dibenzo[b,e][1,4]diazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 15 (R)-1-(3-(6-Oxo-11,12-dihydro-5H-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;
- (R)-1-(3-(10,11-Dihydro-dibenz[b,f][1,4]oxazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 20 (R)-1-(3-(5,6,11,12-Tetrahydro-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;
- 25 (R)-1-(3-(11-Oxo-6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidine-carboxylic acid;

(R)-1-(3-(5-Methyl-dibenz[b,f]azepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

(R)-1-(3-(6,7-Dihydro-5H-dibenz[b,g][1,5]oxazocin-6-yl)-1-propyl)-3-piperidine-
5 carboxylic acid;

(R)-1-(3-(11,12-Dihydro-dibenz[a,e]cycloocten-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

10 or a pharmaceutically acceptable salt thereof.

As used herein, the term "patient" includes any mammal which could benefit from treatment of neurogenic pain or inflammation or insulin resistance in NIDDM. The term particularly refers to a human patient, but is not intended to
15 be so limited.

It has been demonstrated that the novel compounds of formula I inhibit neurogenic inflammation which involves the release of neuropeptides from peripheral and central endings of sensory C-fibres. Experimentally this can be
20 demonstrated in animal models of formalin induced pain or paw oedema (Wheeler and Cowan, Agents Actions 1991, 34, 264-269) in which the novel compounds of formula I exhibit a potent inhibitory effect. Compounds of formula I may be used to treat all painful, hyperalgesic and/or inflammatory conditions in which C-fibers play a pathophysiological role by eliciting neurogenic pain or
25 inflammation, i.e.:


Acutely painful conditions exemplified by migraine, postoperative pain, burns, bruises, post-herpetic pain (Zoster) and pain as it is generally associated with

acute inflammation; chronic, painful and/or inflammatory conditions exemplified by various types of neuropathy (diabetic, post-traumatic, toxic), neuralgia, rheumatoid arthritis, spondylitis, gout, inflammatory bowel disease, prostatitis, cancer pain, chronic headache, coughing, asthma, chronic pancreatitis,
 5 inflammatory skin disease including psoriasis and autoimmune dermatoses, osteoporotic pain.

Further, it has been demonstrated that the compounds of general formula I improves the glucose tolerance in diabetic ob/ob mice and that this may result from the reduced release of CGRP from peripheral nervous endings. Hence the
 10 compounds of general formula I may be used in the treatment of NIDDM as well as aging-associated obesity. Experimentally this has been demonstrated by the subcutaneous administration of glucose into ob/ob mice with or without previous oral treatment with a compound of general formula I.

15

The compounds of formula I may be prepared by the following method:



A compound of formula II wherein R^1 , R^2 , X, Y, p, q and r are as defined above and W is a suitable leaving group such as halogen, p-toluene sulphonate or mesylate may be reacted with an azaheterocyclic compound of formula III wherein R^3 , R^4 , R^5 , m and n are as defined above. This alkylation reaction may
5 be carried out in a solvent such as acetone, dibutylether, 2-butanone, methyl ethyl ketone, ethyl acetate, tetrahydrofuran (THF) or toluene in the presence of a base e.g. potassium carbonate and a catalyst, e.g. an alkali metal iodide at a temperature up to reflux temperature for the solvent used, for e.g. 1 to 120 h. If
10 esters have been prepared in which R^5 is alkoxy, compounds of formula I wherein R^5 is OH may be prepared by hydrolysis of the ester group, preferably at room temperature in a mixture of an aqueous alkali metal hydroxide solution and an alcohol such as methanol or ethanol, for example, for about 0.5 to 6 h.

Compounds of formula II and III may readily be prepared by methods familiar
15 to those skilled in the art.

Under certain circumstances it may be necessary to protect the intermediates used in the above methods e.g. a compound of formula III with suitable protecting groups. The carboxylic acid group can, for example, be esterified.
20 Introduction and removal of such groups is described in "Protective Groups in Organic Chemistry" J.F.W. McOrnie ed. (New York, 1973).

Pharmacological Methods

25 Formalin induced pain or paw oedema

Values for in vivo inhibition of formalin induced pain or oedema for the compounds of the present invention were assessed in mice essentially by the

method of Wheeler-Aceto and Cowan (Agents Action 1991, 34, 265-269).

5 About 20 g NMRI female mice were injected 20 µl 1 % formalin into the left hind paw. The animals were then placed on a heated (31 °C) table, and the pain response was scored. After 1 h they were killed and bled. Left and right hind paws were removed and the weight difference between the paws was used as indication of the oedema response of the formalin injected paw.

Reduced release of CGRP

10 ob/ob female mice, 16 weeks of age, were injected glucose (2g/kg) subcutaneously. At times hereafter blood glucose was determined in tail venous blood by the glucose oxidase method. At the end of the study the animals were decapitated and trunk blood collected. Immunoreactive CGRP was determined in plasma by radio-immuno-assay. Two groups of animals
15 were used. The one group was vehicle treated, whereas the other group received a compound of formula I via drinking water (100 mg/l) for five days before the test.

20 Values for inhibition of formalin induced pain response for some representative compounds are recorded in table 1.

TABLE 1

25 Inhibition of formalin induced pain response at 0.1 mg/kg

Example no.	% Pain inhibition
-------------	-------------------

	1	13
	2	34
	3	36
	4	28
5	6	22

For the above indications the dosage will vary depending on the compound of formula I employed, on the mode of administration and on the therapy desired. However, in general, satisfactory results are obtained with a dosage of from about 0.5 mg to about 1000 mg, preferably from about 1 mg to about 500 mg of compounds of formula I, conveniently given from 1 to 5 times daily, optionally in sustained release form. Usually, dosage forms suitable for oral administration comprise from about 0.5 mg to about 1000 mg, preferably from about 1 mg to about 500 mg of the compounds of formula I admixed with a pharmaceutical carrier or diluent.

The compounds of formula I may be administered in a pharmaceutically acceptable acid addition salt form or where possible as a metal or a lower alkylammonium salt. Such salt forms exhibit approximately the same order of activity as the free base forms.

This invention also relates to pharmaceutical compositions comprising a compound of formula I or a pharmaceutically acceptable salt thereof and, usually, such compositions also contain a pharmaceutical carrier or diluent. The compositions containing the compounds of this invention may be prepared by conventional techniques and appear in conventional forms, for example capsules, tablets, solutions or suspensions.

The pharmaceutical carrier employed may be a conventional solid or liquid carrier. Examples of solid carriers are lactose, terra alba, sucrose, talc, gelatine, agar, pectin, acacia, magnesium stearate and stearic acid. Examples of liquid carriers are syrup, peanut oil, olive oil and water.

5

Similarly, the carrier or diluent may include any time delay material known to the art, such as glyceryl monostearate or glyceryl distearate, alone or mixed with a wax.

- 10 If a solid carrier for oral administration is used, the preparation can be tableted, placed in a hard gelatine capsule in powder or pellet form or it can be in the form of a troche or lozenge. The amount of solid carrier will vary widely but will usually be from about 25 mg to about 1 g. If a liquid carrier is used, the preparation may be in the form of a syrup, emulsion, soft gelatine capsule or
15 sterile injectable liquid such as an aqueous or non-aqueous liquid suspension or solution.

- Generally, the compounds of this invention are dispensed in unit dosage form comprising 50-200 mg of active ingredient in or together with a
20 pharmaceutically acceptable carrier per unit dosage.

The dosage of the compounds according to this invention is 1-500 mg/day, e.g. about 100 mg per dose, when administered to patients, e.g. humans, as a drug.

25

A typical tablet which may be prepared by conventional tableting techniques contains

14

Core:

	Active compound (as free compound or salt thereof)	100 mg
	Colloidal silicon dioxide (Areosil®)	1.5 mg
5	Cellulose, microcryst. (Avicel®)	70 mg
	Modified cellulose gum (Ac-Di-Sol®)	7.5 mg
	Magnesium stearate	

Coating:

10	HPMC	approx.	9 mg
	*Mywacett® 9-40 T	approx.	0.9 mg

*Acylated monoglyceride used as plasticizer for film coating.

- 15 The route of administration may be any route which effectively transports the active compound to the appropriate or desired site of action, such as oral or parenteral e.g. rectal, transdermal, subcutaneous, intranasal, intramuscular, topical, intravenous, intraurethral, ophthalmic solution or an ointment, the oral route being preferred.

20

EXAMPLES

- 25 The process for preparing compounds of formula I and preparations containing them is further illustrated in the following examples, which, however, are not to be construed as limiting.

Hereinafter, TLC is thin layer chromatography, CDCl_3 is deuterio chloroform and DMSO-d_6 is hexadeuterio dimethylsulfoxide. The structures of the


15

compounds are confirmed by either elemental analysis or NMR, where peaks assigned to characteristic protons in the title compounds are presented where appropriate. ^1H NMR shifts (δ_{H}) are given in parts per million (ppm). M.p. is melting point and is given in $^{\circ}\text{C}$ and is not corrected. Column chromatography was carried out using the technique described by W.C. Still et al, J. Org. Chem. (1978), 43, 2923-2925 on Merck silica gel 60 (Art. 9385). Compounds used as starting materials are either known compounds or compounds which can readily be prepared by methods known per se.

10

EXAMPLE 1

(R)-1-(3-(11-Oxo-10,11-dihydro-5H-dibenzo[b,e][1,4]diazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid hydrochloride



15

To a solution of 11-oxo-10,11-dihydro-5H-dibenzo[b,e][1,4]diazepine (12.1 g, 0.058 mol, prepared as described in Synthesis, 1985, 550) in dry N,N-dimethylformamide (150 ml) kept under an atmosphere of nitrogen, sodium hydride (2.8 g, 0.069 mol, 60 % dispersion in oil) was added and the reaction mixture was stirred for 1 hour. 1-Bromo-3-chloropropane (10.9 g, 0.069 mol)

20

was slowly added and the mixture was stirred overnight. The reaction mixture was quenched with water (200 ml) and extracted with diethyl ether (2 x 250 ml). The combined organic extracts were washed with water (3 x 300 ml), dried (MgSO₄) and the solvent was evaporated in vacuo. The solid was washed with
5 heptane (100 ml), filtered off and dried, affording 11.1 g (67 %) of 10-(3-chloro-propyl)-5,10-dihydro-dibenzo[b,e][1,4]diazepin-11-one.

M.p. 127-131°C.

¹H NMR (200 MHz, CDCl₃) δ_H 2.15 (m, 2H), 3.61 (t, 2H), 4.25 (t, 2H), 5.50 (bs, 1H, NH), 6.81 (d, 1H), 6.91 - 7.13 (m, 4H), 7.20 - 7.26 (m, 2H), 7.82 (dd, 1H).
10

A mixture of the above chloride (4.0 g, 0.014 mol), (R)-3-piperidinecarboxylic acid ethyl ester tartrate (8.6 g, 0.028 mol), potassium carbonate (11.7 g, 0.084 mol), potassium iodide (4.6 g, 0.028 mol) and methyl ethyl ketone (100 ml) was
15 heated at reflux temperature for 18 hours under an atmosphere of nitrogen. The cooled reaction mixture was quenched with water (100 ml) and extracted with diethyl ether (100 ml). The organic extract was washed with water (2 x 80 ml), saturated ammonium chloride (80 ml), dried (MgSO₄), filtered and the solvent was evaporated in vacuo. The crude product (7 g) was purified by
20 column chromatography on silica gel (600 ml) using ethyl acetate as eluent. This afforded 3.1 g (55 %) of (R)-1-(3-(11-oxo-10,11-dihydro-5H-dibenzo[b,e]-[1,4]diazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.24 (SiO₂; ethyl acetate).

¹H NMR (200 MHz, CDCl₃) δ_H 1.24 (t, 3H), 1.40 (m, 2H), 1.61 (m, 2H), 1.85 (m, 4H), 2.17 (d, 1H), 2.39 (m, 4H), 2.62 (bd, 1H), 2.82 (bd, 1H), 4.10 (q, 2H), 5.59 (bs, 1H, NH), 6.81 (d, 1H), 6.91 - 7.10 (m, 4H), 7.19 - 7.30 (m, 2H), 7.81
25

(dd, 1H).

The above ethyl ester (2.9 g, 7 mmol) was dissolved in a mixture of ethanol (50 ml) and water (30 ml). Sodium hydroxide (0.34 g, 9 mmol) was added and the
5 reaction mixture was stirred for 40 hours at room temperature. The solvent was evaporated in vacuo, and water (150 ml) was added. The aqueous mixture was washed with diethyl ether (2 x 100 ml), acidified with concentrated hydrochloric acid and washed with dichloromethane (3 x 100 ml). The aqueous phase was evaporated in vacuo. The residue was suspended in a mixture of 2-propanol
10 (10 ml) and diethyl ether (50 ml) and stirred for 60 hours at room temperature. The precipitate was filtered off, washed with diethyl ether and dried in vacuo. Yield 1.57 g (53 %) of the title compound as an amorphous powder.

HPLC retention time = 5.65 min. (5 μ m C18 4 x 250 mm column, eluting with a
15 mixture of water and acetonitril (7:3) containing phosphoric acid and buffered to pH = 3 with triethylamine over 20 minutes at 35 °C).


¹H NMR (400 MHz, DMSO-d₆) δ , 1.27 (m, 1H), 1.50 (d, 1H), 1.59 - 1.78 (m, 5H), 1.91 (m, 1H), 2.22 (t, 2H), 2.57 (d, 1H), 2.85 (d, 1H), 3.38 (m, 1H), 4.00 (t, 2H), 6.94 (t, 1H), 7.10 (m, 3H), 7.20 (m, 1H), 7.32 (m, 2H), 7.59 (d, 1H), 8.02
20 (s, 1H, NH).

By a similar procedure as described in Example 1 the following compound has been prepared:

EXAMPLE 2

(R)-1-(3-(6-Oxo-11,12-dihydro-5H-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid hydrochloride

5



The title compound was prepared by a similar procedure as described in Example 1.

10

HPLC retention time = 7.84 min (5 μ m C18 4 x 250 mm column, eluting for 20 minutes at 30 °C with a mixture of water and acetonitril (7:3) containing phosphoric acid and buffered to pH = 3 with triethylamine).

15


M.p. 223 - 225°C.

MS(EI) 392.2 (M^+ -HCl, 10 %)

20

EXAMPLE 3

(R)-1-(3-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidine-carboxylic acid hydrochloride



5

In a 100 ml round bottom flask equipped with magnetically stirring, thermometer, reflux condenser and addition funnel, 6,11-dihydro-5H-dibenz[b,e]azepine (1.0 g, 5.1 mmol, prepared in a similar way as described in Coll. Czech. Chem. Commun., **23**, 1958, 1330) was dissolved in dry toluene (25 ml). 3-Chloropropionyl chloride (0.78 g, 6.1 mmol) was added slowly. When addition was complete, the reaction mixture was heated at 95 °C for 30 minutes and then allowed to cool to room temperature. With stirring, 0.2 N sodium hydroxide (2.5 ml) was added. More toluene (50 ml) was added and the phases were separated. The organic phase was washed with 0.2 N sodium hydroxide (10 ml) until pH > 10, and then with water (3 x 10 ml) and brine (10 ml). After drying (MgSO₄), the solvent was evaporated in vacuo affording 3-chloro-1-(6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propanone as an oil which was obtained in quantitative yield and used for further reactions without purification.

20

A 1.0 M tetrahydrofuran solution of lithium aluminum hydride (9.8 ml) was

introduced under a nitrogen atmosphere to a dry 100 ml three-necked roundbottom flask using a syringe. Concentrated sulphuric acid (0.264 ml) was cautiously added under cooling on an ice bath, and the resulting solution was stirred at room temperature for 1.5 hour. The above amide was dissolved in dry
5 tetrahydrofuran (9.8 ml) and slowly added. The reaction mixture was stirred overnight. The reaction was quenched by addition of water (0.4 ml), 4 N sodium hydroxide (0.4 ml) and water (1.2 ml), successively. The resulting mixture was filtered (hyflo) and the filtercake was washed with diethyl ether and ethyl acetate. The solvent was evaporated affording 5-(3-chloropropyl)-6,11-
10 dihydro-5H-dibenzo[b,e]azepine, which was used for further reaction without purification.

A solution of potassium iodide (4.5 g, 0.027 mol) in methyl ethyl ketone (75 ml) was heated at reflux temperature for 1 hour. A solution of the above chloride
15 (1.2 g, 4.2 mmol) in methyl ethyl ketone (25 ml) was added and the reaction mixture was heated at reflux temperature for 2.75 hours. (R)-3-Piperidinecarboxylic acid ethyl ester tartrate (1.87 g, 6.2 mmol) and potassium carbonate (1.44 g, 0.01 mol) were added and the mixture was heated at reflux temperature for 24 hours, and left stirring at room temperature for 48 hours.
20 After filtration on filter aid (celite), the solvent was removed by evaporation. The residue was purified by column chromatography on silica gel (125 g) using a mixture of heptane and ethyl acetate (1:1) as eluent. This afforded 0.7 g (42 %) of (R)-1-(3-(6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester as an oil.

25


TLC: R_f = 0.19 (SiO_2 : heptane/ethyl acetate = 1:1).

- The above ester (0.66 g, 1.7 mmol) was dissolved in ethanol (10 ml) and a solution of sodium hydroxide (0.25 g) in water (2 ml) was added. The mixture was stirred at room temperature for 2 hours and concentrated hydrochloric acid (0.8 ml) was added. Dichloromethane (100 ml) was added, followed by water (50 ml) and the phases were separated. The organic phase was dried (MgSO_4) and the solvent evaporated in vacuo. The residue was stripped with acetone, suspended in isopropyl acetate and filtered off. This afforded after drying, 0.5 g (74 %) of the title compound as an amorphous solid.
- 10 HPLC retention time = 20.41 minutes (5 μm C18 4 x 250 mm column, eluting with a 20-80 % gradient of 0.1 % trifluoroacetic acid/acetonitrile and 0.1 % trifluoroacetic acid/water over 30 minutes at 35 °C).
Calculated for $\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_2$, HCl, 0.25 H_2O :
C, 68.14 %; H, 7.28 %; N, 6.91 %; Found:
15 C, 68.08 %; H, 7.44 %; N, 6.61 %.

EXAMPLE 4

20

(R)-1-(3-(6,11-Dioxo-6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidine-carboxylic acid hydrochloride



To a solution of 5H-dibenz[b,e]azepine-6,11-dione (3.0 g, 0.013 mol) in dry N,N-dimethylformamide (25 ml) kept under an atmosphere of nitrogen, sodium hydride (0.7 g, 0.027 mol, 60 % dispersion in mineral oil) was added in portions and the reaction mixture was stirred for 1.5 hour. 1-Bromo-3-chloropropane (5.0 g, 0.031 mol) dissolved in N,N-dimethylformamide was slowly added and the mixture was stirred overnight. Ammonium chloride (2.0 g, 0.04 mol) was added and stirring was continued for 30 minutes. The solution was poured onto water (300 ml) and the mixture was extracted with dichloromethane (2 x 200 ml). The combined organic extracts were dried (MgSO₄), filtered and the solvent was evaporated in vacuo. The residue was purified by column chromatography on silica gel (150 g) using a mixture of heptane and ethyl acetate (4:1) as eluent to give 3.1 g (79 %) of 5-(3-chloropropyl)-5H-dibenz[b,e]azepine-6,11-dione as an oil.

TLC: R_f = 0.48 (SiO₂: heptane/ethyl acetate = 1:1).

A solution of potassium iodide (10.0 g, 0.06 mol) in methyl ethyl ketone (180 ml) was heated at reflux temperature for 1 hour. A solution of the above chloride (3.09 g, 0.01 mol) in methyl ethyl ketone (20 ml) was added and the reaction mixture was heated at reflux temperature for 2 hours. After cooling to

about 60 °C, (R)-3-piperidinecarboxylic acid ethyl ester tartrate (4.50 g, 0.015 mol) and potassium carbonate (3.46 g, 0.025 mol) were added and the mixture was heated at reflux temperature for 48 hours. After cooling and filtration on filter aid (hyflo) the solvent was removed by evaporation. The residue was
5 purified by column chromatography on silica gel (125 g) using first a mixture of heptane and ethyl acetate (1:1) to eluate the front running fractions and then with a mixture of heptane and ethyl acetate (2:3) as eluent. This afforded 1.21 g (29 %) of (R)-1-(3-(6,11-dioxo-6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester as an oil.

10

TLC: R_f = 0.05 (SiO₂: heptane/ethyl acetate = 2:3).

To the above ester (0.60 g, 1.4 mmol) in ethanol (10 ml), a solution of sodium hydroxide (0.35 g) in water (2 ml) was added. The mixture was stirred at room
15 temperature for 2 hours and then concentrated hydrochloric acid (1 ml) was added. Dichloromethane (100 ml) was added, followed by water (50 ml) and the phases were separated. The organic phase was dried (MgSO₄) and the solvent was evaporated in vacuo. Subsequent re-evaporation with acetone and addition of isopropyl acetate afforded, after filtration and drying 0.2 g (23 %) of
20 the title compound as a solid.

HPLC retention time = 16.76 minutes (5 µm C18 4 x 250 mm column, eluting with a 20-80 % gradient of 0.1 % trifluoroacetic acid/acetonitrile and 0.1 % trifluoroacetic acid/water over 30 minutes at 35 °C).

25


¹H NMR (400 MHz, DMSO-d₆) δ_H 1.35 (m, 1H), 1.50 (m, 1H), 1.61 (m, 1H), 1.80 (m, 3H), 2.19 (m, 1H), 2.33 (m, 1H), 2.54 (m, 3H), 2.78 (d, 1H), 2.93 (d,

1H), 3.39 (m, 2H), 4.50 (bs, 1H), 6.40 (t, 1H), 6.81 (d, 1H), 6.93 (d, 1H), 7.27-7.34 (m, 2H), 7.53 (t, 1H), 7.60 (t, 1H), 7.94 (d, 1H), 8.70 (bs, 1H).

5

EXAMPLE 5

10 (R)-1-(3-(10,11-Dihydro-dibenz[b,f][1,4]oxazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid hydrochloride



15 In a dry 250 ml three-necked flask equipped with reflux condenser and thermometer, lithium aluminum hydride (1.8 g, 0.047 mol) was suspended in dry diethyl ether (75 ml) under a nitrogen atmosphere. Cautiously, 10,11-dihydrodibenz[b,f][1,4]oxazepin-11-one (5.0 g, 0.024 mol) was added in portions. The mixture was heated at reflux temperature for 5 hours, cooled to room temperature and left stirring overnight. The reaction mixture was
20 quenched by cautious addition of water (1.8 ml), 4 N sodium hydroxide (1.8 ml) and finally water (5.4 ml). Diethyl ether and toluene were added and the mixture was filtered. The filtercake was washed with diethyl ether, toluene and

ethyl acetate, successively. The filtrate was evaporated until dryness, affording 4.55 g (96 %) of 10,11-dihydrodibenz[b,f][1,4]oxazepine.

5 The above tricycle (4.0 g, 0.02 mol) was dissolved in toluene (100 ml) and 3-chloropropionyl chloride (3.12 g, 0.025 mol) was added. The reaction mixture was heated at 95 °C for 30 minutes and left stirring for 1 hour. 0.2 N Sodium hydroxide (10 ml) was added and the phases were separated. Toluene (200 ml) was added and the organic phase was washed with 0.2 N sodium hydroxide (50 ml), water (3 x 50 ml) and brine (50 ml). The organic phase was
10 dried (MgSO₄), filtered and the solvent evaporated. This afforded 3-chloro-1-(10,11-dihydro-dibenz[b,f][1,4]oxazepin-10-yl)-1-propanone in quantitative yield.

A 1.0 M tetrahydrofuran solution of lithium aluminium hydride (38.4 ml, 0.0384 mol) was introduced into a dry 500 ml three-necked flask. Cautiously,
15 concentrated sulphuric acid (1.84 ml, 0.019 mol) was added dropwise upon cooling on an icebath and then the mixture was stirred for 1.5 hour at room temperature. A solution of the above chloride in dry tetrahydrofuran (38.4 ml) was added dropwise and the reaction mixture was stirred for 1.5 hour. The reaction mixture was subsequently quenched by cautious addition of water (1.5
20 ml), 4 N sodium hydroxide (1.5 ml) and finally water (4.5 ml), and the resulting suspension was left stirring overnight. The precipitated salt was filtered off and washed with diethyl ether, toluene and ethyl acetate. The filtrate was evaporated until dryness to give 4.8 g of crude 10-(3-chloropropyl)-10,11-dihydro-dibenz[b,f][1,4]oxazepine.

25

A solution of potassium iodide (16.0 g, 0.1 mol) in methyl ethyl ketone (180 ml) was heated at reflux temperature for 1 hour. A solution of the above chloride

(3.14 g, 0.012 mol) in methyl ethyl ketone (20 ml) was added and the reaction mixture was heated at reflux temperature for 2.5 hours. (R)-3-Piperidinecarboxylic acid ethyl ester tartrate (7.2 g, 0.024 mol) and potassium carbonate (5.53 g, 0.04 mol) were added and the mixture was heated at reflux temperature for 72 hours. After cooling and filtration on filter aid (celite) the solvent was removed by evaporation. The residue was purified by column chromatography on silica gel (200 g) using a mixture of heptane and ethyl acetate (1:1) as eluent. This afforded 4.5 g (99 %) of (R)-1-(3-(10,11-dihydro-dibenz[b,f][1,4]oxazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.18 (SiO_2 : heptane/ethyl acetate = 1:1).


The above ester (1.0 g, 0.0025 mol) was dissolved in ethanol (15 ml) and a solution of sodium hydroxide (0.38 g) in water (3 ml) was added. The mixture was stirred at room temperature for 2 hours and concentrated hydrochloric acid (1.2 ml) was added. Dichloromethane (150 ml) was added, followed by water (50 ml) and the phases were separated. The organic phase was dried (MgSO_4) and the solvent evaporated in vacuo. Subsequent reevaporation with acetone and addition of a small amount of acetone followed by isopropyl acetate afforded, after filtration and drying 0.2 g (21 %) of the title compound as a solid.

M.p. 185-188 °C.

25 Calculated for $\text{C}_{22}\text{H}_{26}\text{N}_2\text{O}_3 \cdot \text{HCl} \cdot 0.75 \text{H}_2\text{O}$:
C, 63.46 %; H, 6.85 %; N, 6.73 %; Found:
C, 63.06 %; H, 6.74 %; N, 6.47 %.

EXAMPLE 6

5 (R)-1-(3-(5,6,11,12-Tetrahydro-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid dihydrochloride



To a solution of 5,6,11,12-tetrahydro-dibenz[b,f]azocine hydrochloride (7.0 g, 0.029 mol) in dry tetrahydrofuran (100 ml) kept under an atmosphere of nitrogen, n-butyl lithium (22.2 ml, 0.06 mol, 2.7 M in n-hexane) was added dropwise at 0 °C. The reaction mixture was stirred at room temperature for 0.75 hour. A solution of 1-bromo-3-chloropropane (5.4 g, 0.034 mol) in dry tetrahydrofuran (100 ml) was added dropwise at 0 °C and the reaction mixture was stirred at room temperature for 4 hours. The reaction mixture was quenched with water (100 ml) and extracted with diethyl ether (100 ml). The organic extract was washed with brine (80 ml), dried (MgSO₄), filtered, and the solvent was evaporated in vacuo. Yield 7.7 g of crude 5-(3-chloropropyl)-5,6,11,12-tetrahydro-dibenz[b,f]azocine.

20 A mixture of the above chloride (4.7 g, 0.027 mol), (R)-3-piperidinecarboxylic

acid ethyl ester tartrate (16.6 g, 0.054 mol), dry potassium carbonate (22.3 g, 0.162 mol), potassium iodide (8.9 g, 0.054 mmol) and methyl ethyl ketone (250 ml) was heated at reflux temperature for 66 hours under an atmosphere of nitrogen. The cooled reaction mixture was quenched with water (250 ml) and
5 extracted with diethyl ether (250 ml). The organic extract was washed with water (2 x 80 ml), brine (80 ml), dried (MgSO₄), filtered and the solvent was evaporated in vacuo. The crude product (12.4 g) was purified by column chromatography on silica gel (800 ml) using a mixture of ethyl acetate and heptane (1:2) as eluent. This afforded 3.3 g (30 %) of (R)-1-(3-(5,6,11,12-
10 tetrahydro-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.35 (SiO₂; ethyl acetate/heptane = 1:1).

15 The above ethyl ester (3.0 g, 7.38 mmol) was dissolved in a mixture of ethanol (30 ml) and water (30 ml). Sodium hydroxide (0.35 g, 8.85 mmol) was added and the reaction mixture was stirred for 18 hours at room temperature. The solvent was evaporated in vacuo and water (100 ml) was added. The aqueous mixture was washed with diethyl ether (2 x 100 ml) and acidified with
20 concentrated hydrochloric acid. The aqueous phase was evaporated in vacuo and the residue was suspended in 2-propanol (25 ml) and filtered. The filtrate was evaporated in vacuo and the residue was dissolved in acetone (20 ml) and evaporated in vacuo. The residue was dissolved in dichloromethane (100 ml) and filtered. The filtrate was evaporated in vacuo and the residue was
25 suspended in a mixture of acetone (10 ml) and diethyl ether (5 ml) and stirred for 18 hours at room temperature. The precipitate was filtered off, washed with diethyl ether and dried to afford 1.1 g (36 %) of the title compound as a solid.

M.p. 200 - 203 °C

Calculated for $C_{24}H_{30}N_2O_2 \cdot 2HCl, H_2O$:


C, 61.40 %; H, 7.30 %; N, 5.97 %; Found:

C, 61.26 %; H, 7.60 %; N, 5.85 %.

5

EXAMPLE 7

(R)-1-(3-(11-Oxo-6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidine-
10 carboxylic acid hydrochloride



To a mixture of 5H-dibenz[b,e]azepine-6,11-dione (5.0 g, 22.4 mmol), ethylene
15 glycol (12.5 ml, 0.224 mol) and nitromethane (60 ml) in dry toluene (100 ml)
kept under an atmosphere of nitrogen, triflic acid (0.4 ml, 4.5 mmol) was added
dropwise. The reaction mixture was heated at reflux temperature for 3 days
with a water separator. The cooled reaction mixture was quenched with water
(100 ml) and extracted with ethyl acetate (80 ml). The organic extract was
20 washed with water (3 x 80 ml), dried ($MgSO_4$), filtered and the solvent was
evaporated in vacuo. This afforded 5.16 g (86 %) of 5H-dibenz[b,e]azepin-

6,11-dione 11-ethylene ketal as a solid.

TLC: R_f = 0.32 (SiO_2 : heptane/ethyl acetate = 1:1).

^1H NMR (200 MHz, $\text{DMSO}-d_6$) δ , 3.75 (m, 2H), 4.17 (t, 2H), 7.06-7.80 (m, 8H),
5 10.58 (s, 1H).

To a solution of the above ketal (4.5 g, 0.0168 mol) in dry N,N-dimethyl-
formamide (75 ml) kept under an atmosphere of nitrogen, sodium hydride (0.8
g, 0.02 mol, 60 % dispersion in mineral oil) was added in portions and the
10 reaction mixture was stirred for 0.5 hour. A solution of 2-(3-bromo-1-
propyloxy)tetrahydro-2H-pyran (4.5 g, 0.02 mol) in N,N-dimethylformamide was
added slowly and the reaction mixture was stirred for 6 days. The reaction
mixture was poured into water (100 ml) and the mixture was extracted with
diethyl ether (2 x 100 ml). The combined organic extracts were washed with
15 saturated ammonium chloride (3 x 100 ml), dried (MgSO_4), filtered and the
solvent was evaporated in vacuo. This afforded a residue (6.7 g) which was
purified by column chromatography on silica gel (800 ml) using a mixture of
heptane and ethyl acetate (2:1) as eluent. This afforded 4.4 g (64 %) of 5-(3-
(tetrahydro-2H-pyran-2-yloxy)-1-propyl)-5H-dibenz[b,e]azepin-6,11-dione 11-
20 ethylene ketal as an oil.

TLC: R_f = 0.39 (SiO_2 : heptane/ethyl acetate = 1:1).

In a dry 100 ml three-necked flask equipped with reflux condenser and
25 thermometer, lithium aluminum hydride (0.4 g; 10.5 mmol) was suspended in
dry diethyl ether (50 ml) under a nitrogen atmosphere. Cautiously, a solution of
5-(3-(tetrahydro-2H-pyran-2-yloxy)-1-propyl)-5H-dibenz[b,e]azepin-6,11-dione 11-

ethylene ketal (3.9 g) in diethyl ether (30 ml) and tetrahydrofuran (20 ml) was added dropwise. The mixture was heated at reflux temperature for 3 hours, cooled to room temperature and quenched by cautious addition of water (0.5 ml) and 50 % sodium hydroxide (0.5 ml), and filtered. The filtrate was
5 evaporated until dryness affording 3.9 g of an oil which was purified by column chromatography on silica gel (600 ml) using a mixture of heptane and ethyl acetate (4:1) as eluent. This afforded 1.9 g (50 %) of 5-(3-tetrahydro-2H-pyran-2-yloxy)-1-propyl)-5,6-dihydro-dibenz[b,e]azepin-11-one ethylene ketal as an oil.

10 TLC: R_f = 0.43 (SiO_2 : heptane/ethyl acetate = 2:1).

The above ketal (1.8 g, 4.55 mmol) was dissolved in 85 % ethanol (20 ml) and concentrated hydrochloric acid (1 ml) was added. The reaction mixture was heated at reflux temperature for 3.5 hours. Water (100 ml) was added and the
15 mixture was extracted with diethyl ether (2 x 100 ml). The combined organic extracts were washed with water (100 ml) and brine (100 ml), dried (MgSO_4) and filtered. The solvent was evaporated in vacuo to give 5-(3-hydroxy-1-propyl)-5,6-dihydro-5H-dibenz[b,e]azepin-11-one in quantitative yield.

20 M.p. 113 - 115 °C.

To a mixture of the above alcohol (1.0 g, 3.7 mmol) and triethylamine (1 ml) in dichloromethane (30 ml), methanesulfonyl chloride (0.64 g, 5.61 mmol) dissolved in dichloromethane (10 ml) was added dropwise at 15 °C. The
25 reaction mixture was stirred at room temperature for 2 hours and quenched with water. The organic phase was separated, dried (MgSO_4), filtered and evaporated in vacuo. This afforded methanesulfonic acid 3-(11-oxo-5,6-dihydro-

dibenz[b,e]azepin-5-yl)-1-propyl ester as an oil.

TLC: R_f = 0.22 (SiO_2 : heptane/ethyl acetate = 1:1).

- 5 A mixture of the above methanesulfonate (1.3 g, 3.8 mmol), (R)-3-piperidinecarboxylic acid ethyl ester tartrate (2.3 g, 7.5 mmol), dry potassium carbonate (3.1 g, 23 mmol), and methyl ethyl ketone (50 ml) was heated at reflux temperature for 18 hours under an atmosphere of nitrogen. Potassium iodide (0.2 g, 1.2 mmol) was added and the mixture was heated at reflux
- 10 temperature for 7 hours. After cooling, the reaction mixture was quenched with water (50 ml) and extracted with ethyl acetate (50 ml). The organic extract was washed with water (2 x 80 ml), brine (80 ml), dried (MgSO_4), filtered and the solvent was evaporated in vacuo. The crude product (1.4 g) was purified by column chromatography on silica gel (400 ml) using first a mixture of ethyl
- 15 acetate and heptane (1:3) and then a mixture of ethyl acetate and triethylamine (96:4) as eluents. This afforded 0.9 g (59 %) of (R)-1-(3-(11-oxo-5,6-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester as an oil.

- 20 TLC: R_f = 0.12 (SiO_2 : ethyl acetate/heptane = 2:1).

- To the above ester (0.8 g, 2.0 mmol) dissolved in ethanol (25 ml) a solution of sodium hydroxide (94 mg, 2.4 mmol) in water (15 ml) was added and the mixture was stirred at room temperature for 18 hours. The solvent was
- 25 evaporated and water (75 ml) was added to the residue. The aqueous phase was washed with diethyl ether (2 x 75 ml), acidified to pH 1 with concentrated hydrochloric acid and washed with dichloromethane (3 x 100 ml). The aqueous

phase was evaporated in vacuo and dichloromethane (400 ml) was added to the residue. The mixture was stirred for 18 hours, dried (MgSO_4), filtered and evaporated in vacuo. The residue was suspended in acetone (20 ml) and stirred for 18 hours. The solid was filtered off, washed with acetone and dried.

5 This afforded 0.31 g (38 %) of the title compound as a solid.

M.p. 221-223 °C


Calculated for $\text{C}_{23}\text{H}_{28}\text{N}_2\text{O}_3$, HCl, $0.5\text{H}_2\text{O}$:

C, 65.16 %; H, 6.66 %; N, 6.61 %; Found:

10 C, 65.05 %; H, 6.73 %; N, 6.42 %.

EXAMPLE 8

15 (R)-1-(3-(5-Methyl-dibenz[b,f]azepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid hydrochloride



20 To a solution of 10-methoxy-5H-dibenz[b,f]azepine (40.7 g, 0.182 mol, prepared as described in Swiss pat. 389,619) in dry tetrahydrofuran (400 ml) kept under

an atmosphere of nitrogen, n-butyl lithium (80 ml, 0.216 mol, 2.7 M in n-hexane) was added dropwise at -30 °C. The reaction mixture was stirred 0.5 hour. Iodomethane (13.3 ml, 0.214 mol) dissolved in dry tetrahydrofuran (50 ml) was added dropwise at -20 °C. The reaction mixture was allowed to reach 0 °C
5 quenched with water (300 ml) and extracted with diethyl ether (2 x 250 ml). The combined organic extracts were washed with water (500 ml), saturated brine (100 ml) and dried (MgSO_4), filtered and the solvent evaporated in vacuo. Yield 43.5 g (100 %) of 5-methyl-10-methoxy-dibenz[b,f]azepine.

- 10 5-Methyl-10-methoxydibenz[b,f]azepine (43 g, 0.181 mol) was dissolved in 2 N hydrochloric acid (800 ml) and heated at reflux temperature for 1.5 hour. The cooled reaction mixture was extracted with diethyl ether (2 x 400 ml). The combined organic extracts were washed with water (500 ml), saturated brine (100 ml) and dried (MgSO_4), filtered and the solvent was evaporated in vacuo.
15 The crude product (39.5 g) was purified by column chromatography on silica gel (2 x 900 ml) using a mixture of ethyl acetate and heptane (10:1) as eluent. This afforded 21.5 g (53 %) of 10,11-dihydro-5-methyl-dibenz[b,f]azepin-10-one.

M.p. 92 - 93 °C

20

- To an ethylmagnesium bromide solution (prepared from magnesium (2.3 g, 0.094 mol) and ethylbromide (8.1 ml, 0.103 mol) and dry tetrahydrofuran (50 ml)) cooled on an icebath, a solution of 3-chloro-1-propanol (7.9 ml, 0.094 mol) in dry tetrahydrofuran (50 ml) was added dropwise with stirring. After the
25 addition was complete, the mixture was stirred for 0.5 hour. Magnesium (2.3 g, 0.094 mol) and an iodine crystal were added. Stirring was continued at reflux temperature for 1.5 hour. To this mixture a solution of 10,11-dihydro-5-methyl-

dibenz[b,f]azepin-10-one (7.0 g, 0.031 mol) in dry tetrahydrofuran (75 ml) was added dropwise and the reaction mixture was stirred at reflux temperature for 2 hours. The cooled reaction mixture was quenched with saturated ammonium chloride (50 ml), water (50 ml) and extracted with ethyl acetate (2 x 100 ml).

- 5 The combined organic extracts were washed with water (2 x 100 ml), saturated brine (100 ml) and dried (MgSO_4), filtered and the solvent was evaporated in vacuo. The crude product (7 g) was purified by column chromatography on silica gel (700 ml) using a mixture of ethyl acetate and heptane (1:2) as eluent. Yield 0.9 g (11 %) of 3-(5-methyl-dibenz[b,f]azepin-10-yl)-1-propanol.

10

^1H NMR (200 MHz, CDCl_3) δ , 1.90 (bs, 1H), 2.09 (dp, 2H), 3.32 (s, 3H), 3.88 (t, 2H), 4.16 (t, 2H), 6.04 (s, 1H), 6.91-7.34 (m, 7H), 7.45 (dd, 1H).

- 15 To a solution of 3-(5-methyl-dibenz[b,f]azepin-10-yl)-1-propanol (0.8 g, 3.01 mmol) and triethylamine (0.8 ml) in dichloromethane (25 ml), methanesulfonyl chloride (0.5 g, 4.52 mmol) dissolved in dichloromethane (5 ml) was added dropwise at 15 °C. The reaction mixture was stirred at room temperature for 2 hours and quenched with water (25 ml). The organic phase was separated and dried (MgSO_4), filtered and evaporated in vacuo. This afforded 1.0 g (97 %) of
- 20 methanesulfonic acid 3-(5-methyl-dibenz[b,f]azepin-10-yl)-1-propyl ester as an oil.

TLC: R_f = 0.49 (SiO_2 ; heptane/ethyl acetate = 1:1).

- 25 A mixture of the above methanesulfonate (1.0 g, 2.91 mmol), (R)-3-piperidinecarboxylic acid ethyl ester tartrate (1.8 g, 5.82 mmol), dry potassium carbonate (2.4 g, 17.5 mmol), and methyl ethyl ketone (50 ml) was heated at

reflux temperature for 18 hours under an atmosphere of nitrogen. Potassium iodide (0.2 g, 1.205 mmol) was added and the mixture was heated at reflux for an additional 18 hours. The cooled reaction mixture was quenched with water (50 ml) and extracted with ethyl acetate (50 ml). The organic phase was
5 washed with water (2 x 80 ml), saturated brine (80 ml), dried (MgSO_4), filtered and the solvent evaporated in vacuo. The crude product (1.3 g) was purified by column chromatography on silica gel (500 ml) using a mixture of ethyl acetate and heptane (2:1) and later on a mixture of ethyl acetate and triethylamine (96:4) as eluents. This afforded 0.87 g (74 %) of (R)-1-(3-(5-methyl-
10 dibenz[b,f]azepin-10-yl)-1-propyl)-3-piperidine-carboxylic acid ethyl ester as an oil.

TLC: R_f = 0.25 (SiO_2 ; ethyl acetate/heptane = 2:1).

15 The above ester (0.7 g, 1.73 mmol) was dissolved in a mixture of ethanol (25 ml) and water (15 ml), sodium hydroxide (83 mg, 2.08 mmol) was added and the mixture was stirred at room temperature for 60 hours. The solvent was evaporated in vacuo and the residue dissolved in water (50 ml) and washed with diethyl ether (50 ml). The aqueous phase was acidified with concentrated
20 hydrochloric acid until pH = 1 and extracted with dichloromethane (3 x 100 ml). The combined dichloromethane phases were dried (MgSO_4) and the solvent evaporated in vacuo. The residue was suspended in a mixture of diethyl ether (10 ml) and acetone (10 ml) and stirred at room temperature for 18 hours. The solid was filtered off and washed with diethyl ether and dried. This afforded
25 0.15 g (20 %) of the title compound as a solid.

M.p. 222 - 224 °C

37

Calculated for $C_{24}H_{28}N_2O_2 \cdot HCl \cdot 1.25 H_2O$:

C, 66.19 %; H, 7.29 %; N, 6.43 %; Found:


C, 66.01 %; H, 7.00 %; N, 6.15 %.

5

EXAMPLE 9

(R)-1-(3-(6,7-Dihydro-5H-dibenz[b,g][1,5]oxazocin-6-yl)-1-propyl)-3-piperidine-carboxylic acid dihydrochloride

10



15

3-(6,7-Dihydro-5H-dibenz[b,g][1,5]oxazocin-6-yl)-1-propanol (0.60 g, 2.2 mmol, the compound is described in German Patent, DE 2044508 710519 and prepared in a similar way as described in J. Med. Chem., 11, 1968, 97 for an analogous azocine) was suspended in dry toluene (20 ml). On an ice bath and under a nitrogen atmosphere, triethylamine (0.9 ml) was added, followed by methansulfonyl chloride (0.3 ml) which was added dropwise. The resulting mixture was stirred at room temperature for 2 hours. Water (25 ml) was added followed by toluene (50 ml) and dichloromethane (50 ml). The phases were separated. The aqueous phase was extracted with ethyl acetate (50 ml). The combined organic phases were evaporated in vacuo. Methyl ethyl ketone (75 ml) was added to the residue, followed by (R)-3-piperidinecarboxylic acid ethyl

20

25

ester tartrate (1.00 g, 3.3 mmol), potassium iodide (0.36 g, 2.2 mmol), and potassium carbonate (0.76 g, 5.5 mmol). The mixture was heated at reflux temperature for 216 hours. After filtration on filter aid (hyflo), the solvent was removed by evaporation in vacuo. The residue was purified by column
5 chromatography on silica gel (150 ml) using ethyl acetate as eluent. This afforded 0.33 g (37 %) of (R)-1-(3-(6,7-dihydro-5H-dibenz[b,g][1,5]oxazocin-6-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester as an oil.

TLC: R_f = 0.10 (SiO₂:ethyl acetate).

10 The above ester (0.19 g, 0.46 mmol) was dissolved in 96 % ethanol (20 ml) and a solution of sodium hydroxide (0.57 g) in water (3 ml) was added. The mixture was stirred at room temperature for 2 hours and concentrated hydrochloric acid (0.4 ml) was added. Dichloromethane (100 ml) was added, followed by water (50 ml) and the phases were separated. The aqueous phase
15 was washed with dichloromethane (2 x 75 ml) and evaporated in vacuo. Isopropanol was added to the residue and the mixture was filtered. The filtrate was evaporated in vacuo. The resulting solid was redissolved in isopropanol and isopropyl acetate was added. The precipitate was filtered off. This afforded after drying, 0.16 g (78 %) of the title compound as an amorphous solid. The
20 product was further purified for analysis by washing with first dichloromethane and then acetone.

M.p. amorph


Calculated for C₂₃H₂₈N₂O₃, 2 HCl, 1.1 H₂O:

25 C, 58.31 %; H, 6.80 %; N, 5.91 %; Found:

C, 58.78 %; H, 7.26 %; N, 5.53 %.

EXAMPLE 10

5 (R)-1-(3-(11,12-Dihydro-dibenz[a,e]cycloocten-5-yl)-1-propyl)-3-piperidinecarboxylic acid hydrochloride



- 10 To a solution of sodium hydride (2.7 g, 0.066 mol, 60 % dispersion in oil) in ethanol (200 ml) was added triethyl phosphonoacetate (14.9 g, 0.066 mol) and the reaction mixture was stirred at room temperature for 0.5 hour. To the reaction mixture was added 5,6,11,12-tetrahydro-dibenz[a,e]cycloocten-5-one (12.3 g, 0.055 mol, prepared as described in *J. Am. Chem. Soc.*, 77, 1955, 5078) and the reaction mixture was heated at reflux temperature for 18 hours.
- 15 The solvent was evaporated in vacuo and the residue was suspended in water (200 ml) and extracted with diethyl ether (2 x 200 ml). The combined organic extracts were washed with water (2 x 80 ml), saturated ammonium chloride (80 ml), dried (MgSO₄), filtered and the solvent was evaporated in vacuo. The
- 20 crude product (15.7 g) was purified by column chromatography on silica gel

(800 ml) using a mixture of ethyl acetate and heptane (1:10) as eluent. This afforded 7.7 g (48 %) of 11,12-dihydro-dibenz[a,e]cycloocten-5-acetic acid ethyl ester as an oil.

5 TLC: $R_f = 0.21$ (SiO_2 : ethyl acetate/heptane = 1:10).

To a solution of lithiumaluminum hydride (1.9 g, 0.050 mol) in dry diethyl ether: a solution of the above ester (7.3 g, 0.025 mol) in dry diethyl ether (50 ml) was added. The reaction mixture was stirred at reflux temperature for 1.5 hour. The
10 reaction mixture was cooled on a ice bath and quenched by addition of water (2 ml), 20 % sodium hydroxide (2 ml), and MgSO_4 successively. The resulting mixture was filtered and the filtercake was washed with diethyl ether (2 x 100 ml). The phases were separated and the organic phase was evaporated in vacuo affording 6.3 g (100 %) of 2-(11,12-dihydro-dibenz[a,e]cycloocten-5-yl)-
15 ethanol.

TLC: $R_f = 0.16$ (SiO_2 : ethyl acetate/heptane = 1:4).

To a mixture of the above alcohol (6.0 g, 0.024 mol) and triethylamine (6 ml) in
20 dichloromethane (100 ml), methanesulfonyl chloride (4.1 g, 0.036 mol) dissolved in dichloromethane (25 ml) was added dropwise at 15 °C. The reaction mixture was stirred at room temperature for 2 hours and quenched with water (50 ml). The organic phase was separated, dried (MgSO_4) filtered and evaporated in vacuo. This afforded 7.7 g (98 %) of methanesulfonic acid 2-
25 (11,12-dihydro-dibenz[a,e]cycloocten-5-yl)-1-ethyl ester as an oil.

A mixture of the above methanesulfonate (7.7 g), potassium cyanide (2.0 g,

0.031 mol), and dry dimethylsulfoxide (100 ml) was heated at 60 °C for 1.5 hour under an atmosphere of nitrogen. The cooled reaction mixture was quenched with water (100 ml) and extracted with diethyl ether (2 x 100 ml). The organic extract was washed with water (2 x 80 ml), brine (80 ml), dried
5 (MgSO₄), filtered and the solvent was evaporated in vacuo affording 4.7 g (76 %) of 3-(11,12-dihydro-dibenzo[a,e]cycloocten-5-yl)-propionitrile as an oil.

TLC: R_f = 0.72 (SiO₂: ethyl acetate/heptane = 1:2).

- 10 A mixture of the above propionitrile (3.6 g, 0.014 mol) and 50 % potassium hydroxide was heated at reflux temperature for 42 hours. To the cooled reaction mixture water (300 ml) was added and the mixture was washed with diethyl ether (100 ml). The aqueous phase was acidified to pH = 1 with concentrated hydrochloric acid and extracted with diethyl ether (2 x 200 ml).
- 15 The combined organic extracts were washed with water (100 ml), brine (80 ml), dried (MgSO₄), filtered and the solvent was evaporated in vacuo affording 2.9 g (75 %) of 3-(11,12-dihydro-dibenzo[a,e]cycloocten-5-yl)-1-propionic acid as an oil.
- 20 To a solution of lithiumaluminum hydride (0.7 g, 18 mmol) in dry tetrahydrofuran (30 ml) the above propionic acid (2.6 g, 9.3 mmol) in dry diethyl ether (30 ml) was added. The reaction mixture was stirred at reflux temperature for 1.5 hour. The reaction was cooled on a ice bath and quenched by addition of water (0.7 ml), and 20 % sodium hydroxide (0.7 ml), successively. The
25 resulting mixture was filtered and the filtercake was washed with diethyl ether (2 x 100 ml). The phases were separated and the organic phase was evaporated in vacuo affording 2.6 g (100 %) of 3-(11,12-dihydro-

dibenz[a,e]cycloocten-5-yl)-1-propanol as an oil.

TLC: $R_f = 0.34$ (SiO_2 : ethyl acetate/heptane = 1:2).

- 5 To a mixture of the above alcohol (2.0 g, 7.56 mmol) and triethylamine (2 ml) in dichloromethane (50 ml), methanesulfonyl chloride (1.3 g, 11.4 mmol) dissolved in dichloromethane (10 ml) was added dropwise at 10 °C. The reaction mixture was stirred at room temperature for 2 hours and quenched with water (50 ml). The organic phase was separated, dried (Na_2SO_4), filtered and evaporated in
- 10 vacuo. This afforded 2.5 g of methanesulfonic acid 3-(11,12-dihydro-dibenz-[a,e]cycloocten-5-yl)-1-propyl ester as an oil.

- A mixture of the above methanesulfonate (2.5 g), (R)-3-piperidinecarboxylic acid ethyl ester tartrate (4.6 g, 15.1 mmol), dry potassium carbonate (6.3 g,
- 15 45.4 mmol), and methyl ethyl ketone (100 ml) was heated at reflux temperature for 48 hours under an atmosphere of nitrogen. The cooled reaction mixture was quenched with water (100 ml) and extracted with diethyl ether (100 ml). The organic extract was washed with saturated brine (80 ml), dried (Na_2SO_4), filtered and the solvent evaporated in vacuo. The crude product (3.1 g) was
- 20 purified by column chromatography on silica gel (600 ml) using a mixture of ethyl acetate/heptane (1:2) as eluent. This afforded 1.9 g of crude (R)-1-(3-(11,12-dihydro-dibenz[a,e]cycloocten-5-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester. The ethyl ester was suspended in water (150 ml) and concentrated
- 25 hydrochloric acid was added to pH = 1. The aqueous phase was washed with diethyl ether (100 ml) basified to pH = 8 with 20 % sodium hydroxide and extracted with diethyl ether (2 x 100 ml). The combined organic extracts were washed with brine (80 ml), dried (MgSO_4), filtered and the solvent was

evaporated in vacuo affording 1.1 g (36 %) of (R)-1-(3-(11,12-dihydro-dibenz[a,e]cycloocten-5-yl)-1-propyl)-3-piperidinecarboxylic acid ethyl ester as an oil.

5 TLC: $R_f = 0.42$ (SiO_2 : ethyl acetate/heptane = 1:1).

A solution of sodium hydroxide (120 mg, 3.0 mmol) in water (10 ml) was added to the above ester (1.0 g, 2.5 mmol) dissolved in ethanol (25 ml), and the mixture was stirred at room temperature for 60 hours. The solvent was
10 evaporated in vacuo and to the residue water (100 ml) was added. The aqueous mixture was washed with diethyl ether (2 x 50 ml) acidified to pH = 1 with concentrated hydrochloric acid and extracted with dichloromethane (3 x 75 ml). The combined organic extracts were dried (MgSO_4), filtered and evaporated in vacuo. This afforded 0.67 g (66 %) of the title compound as a
15 solid.


Calculated for $\text{C}_{25}\text{H}_{29}\text{NO}_2 \cdot \text{HCl} \cdot 0.25 \text{H}_2\text{O}$:

C, 72.10 %; H, 7.38 %; N, 3.36 %; Found:

20 C, 71.88 %; H, 7.67 %; N, 3.09 %.

CLAIMS

1. A compound of formula I



wherein R^1 and R^2 independently are hydrogen, halogen, trifluoromethyl, hydroxy, C_{1-6} -alkyl or C_{1-6} -alkoxy; and

15 Y is $-\underline{CH}_2-\underline{N}-CH_2-$, $-\underline{CH}_2-\underline{N}-\underline{CH}_2-$, $-(\underline{C}=\underline{O})-\underline{N}-CH_2-$, $-\underline{CH}_2-\underline{N}-(\underline{C}=\underline{O})-$, $-\underline{CH}=\underline{C}-CH_2-$, $-\underline{CH}_2-\underline{C}=\underline{CH}-$, $-\underline{CH}_2-\underline{CH}-CH_2-$, $-\underline{CH}_2-\underline{CH}-\underline{CH}_2-$, $-\underline{CH}_2-\underline{C}=\underline{CH}-$, $-\underline{CH}=\underline{C}-CH_2-$, $-\underline{O}-\underline{CH}-CH_2-$, $-\underline{CH}_2-\underline{CH}-\underline{O}-$, $-\underline{S}-\underline{CH}-CH_2-$ or $-\underline{CH}_2-\underline{CH}-\underline{S}-$ wherein only the underscored atoms participate in the ring system; and

20 X is $-O-$, $-S-$, $-C(R^6R^7)-$, $-CH_2CH_2-$, $-CH=CH-CH_2-$, $-CH_2-CH=CH-$, $-CH_2-(C=O)-$, $-(C=O)-CH_2-$, $-CH_2CH_2CH_2-$, $-CH=CH-$, $-N(R^8)-(C=O)-$, $-(C=O)-N(R^8)-$, $-O-CH_2-$, $-CH_2-O-$, $-S-CH_2-$, $-CH_2-S-$, $-(C=O)-$, $-N(R^9)-$ or $-(S=O)-$ wherein R^6 , R^7 , R^8 and R^9 independently are hydrogen or C_{1-6} -alkyl; and q is 0 or 1; and

p is 0 or 1; and

25 r is 1, 2 or 3; and

m is 1 or 2; and

n is 1 when m is 1 and n is 0 when m is 2; and

R^3 and R^4 each represents hydrogen or may - when m is 2 - together

represent a bond; and

R⁶ is -OH or C₁₋₆-alkoxy;

or a pharmaceutically acceptable salt thereof.

- 5 2. A compound according to claim 1 wherein R¹ and R² independently represent hydrogen, halogen or C₁₋₆-alkoxy.
3. A compound according to the previous claim wherein R¹ and R² are hydrogen.
- 10 4. A compound according to any one of the preceding claims wherein Y is -CH₂-N-CH₂-, -CH₂-N-CH₂-, -(C=O)-N-CH₂-, -CH₂-N-(C=O)-, -CH₂-C=CH-, -CH=C-CH₂-, -CH₂-C=CH- or -CH=C-CH₂-.
- 15 5. A compound according to the previous claim wherein Y is -CH₂-N-CH₂-, -CH₂-N-CH₂-, -(C=O)-N-CH₂-, -CH₂-N-(C=O)- or -CH₂-C=CH-.
6. A compound according to any one of the preceding claims wherein X represents -O-, -S-, -CH₂-, -CH₂CH₂-, -CH=CH-CH₂-, -CH₂-CH=CH-, -CH₂CH₂CH₂-, -CH=CH-, -O-CH₂-, -CH₂-O-, -S-CH₂-, -CH₂-S-, -(C=O)- or -N(R⁹)-, wherein R⁹ is hydrogen or C₁₋₆-alkyl.
- 20 7. A compound according to the previous claim wherein X is -O-, -CH₂-, -CH₂-CH₂-, -(C=O)-, -NH- or -N(CH₃)-.
- 25 8. A compound according to any one of the preceding claims wherein r is 2.

9. A compound according to any one of the preceding claims wherein m is 2.

10. A compound according to any one of the preceding claims wherein R⁵ is -OH.

11. A compound according to any one of the claims 1 through 10 selected from the following:

10 (R)-1-(3-(6,11-Dioxo-6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

(R)-1-(3-(6,11-Dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

15 (R)-1-(3-(5,11-Dihydro-10H-dibenzo[b,e][1,4]diazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

20 (R)-1-(3-(11H-Dibenzo[b,f][1,4]thiazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

(R)-1-(3-(11H-Dibenz[b,f][1,4]oxazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

25 (R)-1-(3-(11H-Dibenz[b,f][1,4]oxathiepin-11-yl)-1-propyl)-3-piperidinecarboxylic acid;

(R)-1-(3-(11H-Dibenzo[b,e][1,4]dithiepin-11-yl)-1-propyl)-3-piperidinecar-

boxylic acid;

(R)-1-(3-(11H-Dibenz[b,e][1,4]oxathiepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

5

(R)-1-(3-(11,12-Dihydro-10H-dibenz[b,g][1,5]oxazocin-11-yl)-1-propyl)-3-piperidinecarboxylic acid;

10

(R)-1-(3-(11,12-Dihydro-10H-dibenzo[b,g][1,5]thiazocin-11-yl)-1-propyl)-3-piperidinecarboxylic acid;

1-(3-(11,12-Dihydro-6H-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

15

1-(3-(11,12-Dihydro-5H-dibenzo[a,e]cycloocten-5-ylidene)-1-propyl)-3-piperidinecarboxylic acid;

20

1-(3-(6-Oxo-11,12-dihydro-5H-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

1-(3-(7,12-Dihydro-6H-dibenzo[a,d]cycloocten-6-ylidene)-1-propyl)-3-piperidinecarboxylic acid;

25

1-(3-(5-Methyl-5,11-dihydro-dibenz[b,f]azepin-10-ylidene)-1-propyl)-3-piperidinecarboxylic acid;

1-(3-(6-Oxo-5,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

(R)-1-(3-(11-Oxo-10,11-dihydro-5H-dibenzo[b,e][1,4]diazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

5 (R)-1-(3-(6-Oxo-11,12-dihydro-5H-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

(R)-1-(3-(10,11-Dihydro-dibenz[b,f][1,4]oxazepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

10

(R)-1-(3-(5,6,11,12-Tetrahydro-dibenz[b,f]azocin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

15

(R)-1-(3-(11-Oxo-6,11-dihydro-5H-dibenz[b,e]azepin-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

(R)-1-(3-(5-Methyl-dibenz[b,f]azepin-10-yl)-1-propyl)-3-piperidinecarboxylic acid;

20

(R)-1-(3-(6,7-Dihydro-5H-dibenz[b,g][1,5]oxazocin-6-yl)-1-propyl)-3-piperidinecarboxylic acid;

(R)-1-(3-(11,12-Dihydro-dibenz[a,e]cycloocten-5-yl)-1-propyl)-3-piperidinecarboxylic acid;

25

or a pharmaceutically acceptable salt thereof.

12. The use of a compound according to any of the claims 1 through 11 as a medicament.

13. The use of a compound according to any of the claims 1 through 11 for preparing a medicament for the treatment of neurogenic inflammation.

14. The use of a compound according to any of the claims 1 through 11 for preparing a medicament for the treatment of insulin resistance in NIDDM or aging.

15. A pharmaceutical composition comprising as active component a compound according to any of the claims 1 through 11 together with a pharmaceutically carrier or diluent.

16. A pharmaceutical composition suitable for treating neurogenic inflammation comprising an effective amount of a compound according to any one of the claims 1 through 11 together with a pharmaceutically acceptable carrier or diluent.

17. A pharmaceutical composition suitable for treating insulin resistance in NIDDM or aging comprising an effective amount of a compound according to any one of the claims 1 through 11 together with a pharmaceutically acceptable carrier or diluent.

18. The pharmaceutical composition according to claims 15, 16 or 17 comprising between 0.5 mg and 1000 mg of the compound according to any one of the claims 1 through 11 per unit dose.

19. A method of treating neurogenic inflammation in a subject in need of such treatment comprising administering to said subject an effective amount of a compound according to any one of the claims 1 through 11.

5 **20.** A method of treating insulin resistance in NIDDM or aging in a subject in need of such treatment comprising administering to said subject an effective amount of a compound according to any one of the claims 1 through 11.


10 **21.** A method of treating neurogenic inflammation in a subject in need of such treatment comprising administering to said subject a pharmaceutical composition according to claim 16.

15 **22.** A method of treating insulin resistance in NIDDM or aging in a subject in need of such treatment comprising administering to said subject a pharmaceutical composition according to claim 17.

20 **23.** A method of preparing a compound according to claim 1,
CHARACTERIZED in

a) reacting a compound of formula II

25



wherein R^1 , R^2 , X , Y , p , q and r are as defined above and W is a suitable leaving group such as halogen, p-toluene sulphonate or mesylate, with a compound of formula III



wherein R^3 , R^4 , R^5 , m and n are as defined above to form a compound of formula I; or

b) hydrolyzing a compound of formula I, wherein R^5 is C_{1-8} -alkoxy, to form a compound of formula I wherein R^5 is OH.

24. Any novel feature or combination of features described herein.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/DK 96/00138

A. CLASSIFICATION OF SUBJECT MATTER		
IPC6: C07D 401/06, C07D 413/06, C07D 211/60, A61K 31/55, A61K 31/445 According to International Patent Classification (IPC) or to both national classification and IPC		
B. FIELDS SEARCHED		
Minimum documentation searched (classification system followed by classification symbols)		
IPC6: C07D		
Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched		
SE,DK,FI,NO classes as above		
Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)		
CAS-ONLINE		
C. DOCUMENTS CONSIDERED TO BE RELEVANT		
Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	GB 1109847 A (RHONE-POULENC S.A.), 18 April 1968 (18.04.68) --	1-10,12-18, 23
A	US 5354747 A (DONALD W. HANSEN, JR. ET AL), 11 October 1994 (11.10.94) --	1-10,12-18, 23
A	DE 1084267 B (CILAG-CHEMIE AKTIENGESELLSCHAFT), 30 June 1960 (30.06.60) --	1-10,12-18, 23
A	US 3803143 A (SATORU TANAKA ET AL), 9 April 1974 (09.04.74) -- -----	1-10,12-18, 23
<input type="checkbox"/> Further documents are listed in the continuation of Box C. <input checked="" type="checkbox"/> See patent family annex.		
* Special categories of cited documents: "A" document defining the general state of the art which is not considered to be of particular relevance "E" earlier document but published on or after the international filing date "L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified) "O" document referring to an oral disclosure, use, exhibition or other means "P" document published prior to the international filing date but later than the priority date claimed "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention "X" document of particular relevance: the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone "Y" document of particular relevance: the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art "&" document member of the same patent family		
Date of the actual completion of the international search		Date of mailing of the international search report
2 July 1996		05 -08- 1996
Name and mailing address of the ISA/ Swedish Patent Office Box 5055, S-102 42 STOCKHOLM Facsimile No. +46 8 666 02 86		Authorized officer Göran Karlsson Telephone No. +46 8 782 25 00

INTERNATIONAL SEARCH REPORT

International application No.

PCT/DK 96/00138

Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This international search report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☒ Claims Nos.: 11, 19-22
because they relate to subject matter not required to be searched by this Authority, namely:
A method for treatment of the human or animal body by therapy,
see Rule 39.1
2. ☒ Claims Nos.: 24
because they relate to parts of the international application that do not comply with the prescribed requirements to such an extent that no meaningful international search can be carried out, specifically:
Claim 24 does not clearly define the matter for which protection is sought, see Article 6
3. ☐ Claims Nos.:
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

1. ☐ As all required additional search fees were timely paid by the applicant, this international search report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this international search report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☐ No required additional search fees were timely paid by the applicant. Consequently, this international search report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
☐ No protest accompanied the payment of additional search fees.

5776.5

INTERNATIONAL SEARCH REPORT
Information on patent family members

01/04/96

International application No.

PCT/DK 96/00138

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
GB-A- 1109847	18/04/68	BE-A- 664609	00/00/00
		CH-A- 436269	00/00/00
		CH-A- 448071	00/00/00
		DE-A- 1620156	02/01/70
		FR-E- 89653	00/00/00
		FR-E- 89659	00/00/00
		FR-A- 1488211	00/00/00
		LU-A- 48732	02/08/65
		NL-A- 6506574	02/12/65
		SE-B- 312549	21/07/69
		US-A- 3480624	25/11/69
		BR-D- 6570074	00/00/00
US-A- 5354747	11/10/94	AU-A- 7138794	03/01/95
		CA-A- 2165159	22/12/94
		EP-A- 0703908	03/04/96
		US-A- 5461047	24/10/95
		WO-A- 9429286	22/12/94
DE-B- 1084267	30/06/60	NONE	
US-A- 3803143	09/04/74	DE-A- 2044508	19/05/71
		FR-A, A, B 2070678	17/09/71
		GB-A- 1293072	18/10/72